

University of Texas at Tyler

## Scholar Works at UT Tyler

---

Biology Faculty Publications and Presentations

Biology

---

11-1-2008

### **Pseudomonas aeruginosa elastase induces IL-8 production in the lung cells via the epidermal growth factor/extracellular signal-regulated proteins/NF $\kappa$ B pathway**

Ali Azghani

University of Texas at Tyler, aazghani@uttyler.edu

Follow this and additional works at: [https://scholarworks.uttyler.edu/biology\\_fac](https://scholarworks.uttyler.edu/biology_fac)



Part of the [Biology Commons](#)

---

#### **Recommended Citation**

Azghani, Ali, "Pseudomonas aeruginosa elastase induces IL-8 production in the lung cells via the epidermal growth factor/extracellular signal-regulated proteins/NF $\kappa$ B pathway" (2008). *Biology Faculty Publications and Presentations*. Paper 1.

<http://hdl.handle.net/10950/246>

This Article is brought to you for free and open access by the Biology at Scholar Works at UT Tyler. It has been accepted for inclusion in Biology Faculty Publications and Presentations by an authorized administrator of Scholar Works at UT Tyler. For more information, please contact [tgullings@uttyler.edu](mailto:tgullings@uttyler.edu).

**Pseudomonas aeruginosa elastase induces IL-8 production in the lung cells via the epidermal growth factor/extracellular signal-regulated proteins/NFκB pathway**

---

**Ali Azghani**

*Department of Biology, The University of Texas at Tyler, Tyler,*

*Texas, USA*

*Critical Care* 2008, **12(Suppl 5)**:P6 (doi: 10.1186/cc7039)

**Background** The induction of chemokine secretion by fibroblasts is crucial for the migration of leukocytes into the parenchyma of the injured lung. Several bacterial products activate the lung's structural as well as immune cells to produce pro inflammatory cytokines and chemokines. We report that elastase from *Pseudomonas aeruginosa* (PE) evokes IL-8 mRNA expression and protein secretion in nonmalignant culture of human lung fibroblasts by activating the receptor for epidermal growth factor (EGFR) and downstream mitogen-activated protein kinases (MAPK) pathway.

**Methods** We utilized western blot analysis to detect phosphorylation of EGFR and signal transduction intermediates. Northern blot and ELISA analyses were used to determine IL-8 RNA expression and cytokine secretion.

**Results** We found that the enzymatically active PE enhances IL-8mRNA and protein secretion but does not increase IL-10 or TNF expression. PE induces phosphorylation of the EGFR and the extracellular signal-regulated proteins (ERK1/2) of the MAPK pathway. Pretreatment of the cells with neutralizing antibody to EGFR or the EGFR-specific tyrphostin AG1478 markedly attenuated the PE-induced ERK1/2 activation. PE-induced IL-8 expression is also abolished in the presence of the MEK inhibitor U0126, indicating the involvement of ERK1/2 in this process.

**Conclusion** Taken together, the results show PE could modulate lung inflammation by exploiting the EGFR/ERK/NFκB pathway and enhancing IL-8 production by lung fibroblasts.