

University of Texas at Tyler

Scholar Works at UT Tyler

Faculty Posters

Research Posters

9-2014

Comparison of Alcohol-Based Sanitizers Versus Personal Protective Equipment on the Incidence of Hospital-associated Infections

Danice B. Greer

The University of Texas at Tyler, dgreer@uttyler.edu

Colleen Marzilli

University of Texas at Tyler, cmarzilli@uttyler.edu

Follow this and additional works at: https://scholarworks.uttyler.edu/fac_posters



Part of the [Nursing Commons](#)

Recommended Citation

Greer, Danice B. and Marzilli, Colleen, "Comparison of Alcohol-Based Sanitizers Versus Personal Protective Equipment on the Incidence of Hospital-associated Infections" (2014). *Faculty Posters*. Book 13.

<http://hdl.handle.net/10950/381>

This Book is brought to you for free and open access by the Research Posters at Scholar Works at UT Tyler. It has been accepted for inclusion in Faculty Posters by an authorized administrator of Scholar Works at UT Tyler. For more information, please contact tgullings@uttyler.edu.

Comparison of Alcohol-Based Sanitizers Versus Personal Protective Equipment on the Incidence of Hospital-associated Infections

Danice B. Greer, PhD, RN, BC, & Colleen Marzilli, DNP, RN, BC, CNE

The University of Texas at Tyler - School of Nursing & Health Sciences

PICO Question: In healthcare settings what is the effect of alcohol-based sanitizers compared to usual personal protective equipment on the incidence of infections?

POPULATION

The CDC healthcare-associated infection (HAI) prevalence survey provides an updated national estimate of the overall problem of HAIs in U.S. hospitals. Based on a large sample of U.S. acute care hospitals, on any given day, about 1 in 25 hospital patients has at least one HAI.

There were an estimated 722,000 HAIs in U.S. acute care hospitals in 2011. About 75,000 hospital patients with HAIs died during hospitalizations. More than half of all HAIs occurred outside of the intensive care unit.¹

Major Site of Infection	Estimated No.
Pneumonia	157,500
Gastrointestinal Illness	123,100
Urinary Tract Infections	93,300
Primary Bloodstream Infections	71,900
Surgical site infections from any inpatient surgery	157,500
Other types of infections	118,500
Estimated total number of infections in hospitals	721,800

Magill, S.S., Edwards, J.R., Bamberg, W., et al. (2014). Multistate point-prevalence survey of health care-associated infections. *New England Journal of Medicine*, 370(13), 1198-1208. DOI: 10.1056/NEJMoa1306801

INTERVENTION

- Alcohol-based sanitizers
- Chlorhexidine
- Iodines
- Handwashing
- Gloves
- Goggles
- Face Mask
- Gowns
- Shoe Covers



CLINICAL DECISION

- Based on the evidence alcohol-based products are more effective for standard hand hygiene/hand antisepsis, however Concentrations of alcohol-based sanitizers vary.
- The most effective in fighting hospital-associated infections are those with concentrations between 60% and 90%.
- Using a combination of alcohol based Sanitizers and gloves is also recommended in fighting HAIs.

STRENGTH OF EVIDENCE

Level of Evidence

Citation

1, 2, 3, 4, & 7	Centers for Disease Control and Prevention, (October 25, 2002). Guideline for hand hygiene in health-care settings: Recommendations of the healthcare infection control practices advisory committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. <i>MMWR</i> , 51(No. RR-16).
1	Jefferson, T., Del Mar, C.B., Dooley, L., et al. (2011). Physical interventions to interrupt or reduce the spread of respiratory viruses. <i>Cochrane Database of Systematic Reviews</i> , 2011, Issue 7. Art. No.: CD006207.
4	Matsumoto, K., Shigemi, A., Yaji, K., et al. (2012). Reduction in the incidence of MRSA with use of alcohol-based hand rub solutions and gloves. <i>Journal of Infection and Chemotherapy</i> , 18(2), 269-271.
1	Montville, R., Schaffner, D.W. (2011). A meta-analysis of the published literature on the effectiveness of antimicrobial soaps. <i>Journal of Food Protection</i> , 74(11), 1875-1882.
3	Sakamoto, F., Yamada, H., Suzuki, C., et al. (2010). Increased use of alcohol-based hand sanitizers and successful eradication of methicillin-resistant staphylococcus aureus from a neonatal intensive care unit: A multivariate time series analysis. <i>American Journal of Infection Control</i> , 38(7), 529-534.
1	Sroka, S., Gastmeier, P., & Meyer, E. (2010). Impact of alcohol hand-rub use on methicillin-resistant Staphylococcus aureus: An analysis of the literature. <i>Journal of Hospital Infection</i> , 74(3), 204-211.
1	Tanner, J., Swarbrook, S., & Stuart, J. (2008). Surgical hand antisepsis to reduce surgical site infection. <i>Cochrane Database of Systematic Reviews</i> , 2008, Issue 1. Art. No.: CD004288.
2	Vernaz, N., Sax, H., Pittet, D., et al. (2008). Temporal effects of antibiotic use and hand rub consumption on the incidence of MRSA and Clostridium difficile. <i>The Journal of Antimicrobial Chemotherapy</i> , 62(3), 601-607.