5-2017

Triggers and Their Influence on Health Behaviors

Christine S. Gipson

University of Texas at Tyler, cgipson@uttyler.edu

Follow this and additional works at: https://scholarworks.uttyler.edu/nursing_fac

Part of the [Nursing Commons](#)

Recommended Citation

Gipson, Christine S., "Triggers and Their Influence on Health Behaviors" (2017). *Nursing Faculty Publications and Presentations*. Paper 20.

<http://hdl.handle.net/10950/1193>

This Article is brought to you for free and open access by the School of Nursing at Scholar Works at UT Tyler. It has been accepted for inclusion in Nursing Faculty Publications and Presentations by an authorized administrator of Scholar Works at UT Tyler. For more information, please contact tbianchi@uttyler.edu.

ARTICLES AND ESSAYS

Triggers and Their Influence on Health Behaviors

Christine S. Gipson, PhD, RN, CNE

This article provides a conceptual definition of the concept trigger within the context of health behaviors and applies it to the highly significant health issue of obesity. Healthy behaviors are essential to life and happiness, but they do not just happen. They are triggered, and an inner drive keeps them alive. To help patients gain and retain optimal health, nurses must understand the triggers of healthy behaviors. Walker and Avant's (2011) method of concept analysis is used as the basis for defining the concept of trigger. The antecedents, defining attributes, and consequences of trigger are identified. Findings suggest that nurses can play a role in triggering health behavior change through simple motivational efforts.

Keywords: trigger; behavior change; influencing behavior; health behaviors

Once considered to be an age of optimal health, the move from adolescence to young adulthood is now recognized as a critical period for health promotion and prevention of disease (Nelson, Kocos, Lytle, & Perry, 2009). Young adulthood is a time of change that includes multiple stressors. Nelson, Lust, Story, and Ehlinger (2008) described the unique combination of stressors such as financial issues, poor dietary intake, and physical inactivity that greatly impact students' short-term and long-term health. Furthermore, lifestyle factors (Mohindra, Nicklas, O'neil, Yang, & Berenson, 2009), along with social influences and the environment, contribute to the increase in obesity by encouraging or intensifying unhealthy behaviors (Eaton, Kann, Okoro, & Collins, 2007; Jago et al., 2004; Quick, Wall, Larson, Haines, & Neumark-Sztainer, 2013). As students transition from a time of dependence on their parents to a time of independence, there are changes and decisions that must be made on a daily basis.

Obesity is a complex problem with multiple social, environmental, behavioral, and personal factors influencing the behavior of young adults. Weight gain in young adulthood is a multifaceted issue that is not completely understood. Understanding the numerous factors that influence behavior and result in behavior change is necessary to decrease the current obesity epidemic. According to Fogg (2011), there are three elements necessary for behavior change. These elements are motivation, ability, and triggers, and they must all occur at the same time for

Christine S. Gipson, PhD, RN, CNE, is an assistant professor and coordinator of the RN-to-BSN Track in the College of Nursing and Health Sciences at the University of Texas at Tyler.

Once considered
to be an age of
optimal health,
the move from
adolescence
to young
adulthood is now
recognized as a
critical period for
health promotion
and prevention of
disease.

behavior change to occur. The term *trigger* is often interchanged with other words such as prompt, cue, call to action, request, or offer (Fogg, 2011). However, trigger is a concept with unique characteristics that separates it from other terms. This article was written to provide a conceptual definition of the concept trigger and its application, as well as its significance, to the nursing profession.

SIGNIFICANCE OF THE CONCEPT FOR NURSING

The term *trigger* has been used in the nursing literature to describe events that precipitate medical conditions such as migraines, cardiac events, and asthma. For these conditions, triggers are well defined and have been studied at length. Moreover, in many cases, scales have been developed to measure their presence. However, this concept as it relates to behavior change is not as well defined. Chambers and Swanson (2012), in a study about factors associated with successful and unsuccessful weight maintenance, described the need to define a clear trigger point for action to help counteract the tendencies of individuals to put off making a change. Visram, Crosland, and Cording (2009) described triggers to behavior change with the subthemes of intrinsic motivation and extrinsic factors. Furthermore, concerns about health, loss of function or mobility, and cosmetic factors were listed as triggers to action for weight loss. The stigma attached to those who are overweight can serve as a catalyst for behavior change (Visram et al., 2009).

CONCEPT IDENTIFICATION

Used as a noun in the *Merriam-Webster's online dictionary*, *trigger* (n.d.) is defined as "something that causes something else to happen." Different uses of the term *trigger* can be found throughout the literature: Asthma triggers, migraine triggers, trigger tool method, trigger points, and pull the trigger are examples. The concept trigger has been used in criminal justice, medicine, academia, and behavioral science.

Criminal justice and the firearm industry use the word *trigger* to describe the lever on a gun which the user pulls to initiate an action (Trigger, n.d.). This overt, conscious, and irrevocable action provides a clear mental image. There are different triggers that can be installed on firearms which enhance this action. When the trigger is engaged, a bullet explodes through the barrel, creating action. The trigger releases the hammer which strikes the bullet causing it to fire. A safety is located on the gun to keep the trigger from firing inadvertently.

In the medical literature, De Wet and Bowie (2011) used a trigger tool to screen medical records for occurrences that alert reviewers to possible adverse events that had previously been undetected. Detection of possible patient harm occurrences led the reviewer to take action to prevent further harm to the patient. Other studies in the medical literature have used trigger tools to alert providers to potential adverse events (Lim, Melucci, Rizer, Prier, & Weber, 2016; Wong et al., 2015). Peterson, Gaeta, Birkhahn, Fernández, and Mancuso (2012) discussed the various triggers that precipitate asthma symptoms, with importance placed on identifying and self-managing asthma triggers as a cornerstone of treatment. Individuals are often aware of specific factors that precipitate asthma symptoms, so accurate identification of asthma triggers is necessary for adequate management (Peterson et al., 2012). Managing asthma involves the use of pharmaceuticals during times of exacerbation and avoiding triggers when possible. The most common trigger

reported in an analysis of migraine triggers by Andress-Rothrock, King, and Rothrock (2010) was emotional stress, followed by lack of or too much sleep, odors, and skipping meals. To be considered a migraine trigger, exposure to the factor must consistently and promptly produce migraine symptoms. Avoiding migraine triggers was the mainstay of treatment (Andress-Rothrock et al., 2010). Studies reviewed in this discipline point to irritants that include internal and external influences that trigger or set off a chain of events in the human body.

A review of the literature in academia was concerned with determining what factors trigger learning or engagement in learning. O'Toole, Solan, Burkhardt, and Klein (2013) examined the use of a video trigger curriculum to increase the frequency of medical residents' screening for social determinants of health (SDH). In the study, residents were shown videos depicting various clinical scenarios that modeled appropriate and inappropriate techniques for screening. Residents who received the trigger curriculum demonstrated a significant change in screening for some SDH and were more likely to report knowledge and comfort to do so (O'Toole et al., 2013). A study on adult lifelong learning conducted by Kungu and Machtmes (2009) argued that understanding the triggers of participation by adults in their learning can provide insight for defining the concept of lifelong learning. In their study, the authors considered circumstances that triggered adult learners to engage in learning. Triggers, when used in the academic setting, have potential to create action in the learner.

Psychology and behavioral management research use the term *trigger* to describe something, whether it be emotional distress or an external stimulus, that *triggers* another behavior (Dalle Grave, Centis, Marzocchi, El Ghoch, & Marchesini, 2013). Furthermore, motivation for change is imperative to modify unhealthy habits, which further necessitate the identification of triggers that lead to behavior change (Dalle Grave et al., 2013). A main point of emphasis found in this discipline was the importance of stimulus control.

Fogg (2009) developed a behavior model that posits that motivation, ability, and trigger are all needed simultaneously to produce behavior change. He describes three types of triggers: spark, facilitator, and signal, in which the motivation and ability of the individual determine the type of trigger needed to produce behavior change. A spark motivates behavior, a facilitator makes the target behaviors easier, and a signal reminds (Fogg, 2009). Each of these triggers spurs an individual into action depending on the level of motivation and ability.

A goal of this review is an exploration of the concept of trigger. After an extensive review of the literature, the working definition of the term *trigger* for the purposes of this article is "an external or internal spark that generates action within an individual."

DEFINING ATTRIBUTES

Defining attributes are characteristics that are frequently associated with a concept and are found repeatedly in the literature (Walker & Avant, 2011). The following defining attributes of trigger were selected:

- Unique to each individual
- An impulse that results in definitive action
- Internal or external spark

Three elements—
motivation, ability,
and triggers—
must all occur at
the same time for
behavior change
to occur.

A defining attribute of trigger is that it is “unique to each individual.” A common theme throughout the literature review was the individuality of stimuli that produced action in different individuals. Another defining attribute of trigger is “an impulse that results in definitive action.” Chambers and Swanson (2012) describe an alarm signal that elicits action in an individual. This trigger signal is that internal or external spark that tells an individual to take action.

ANTECEDENTS AND CONSEQUENCES

Antecedents are actions that must happen before the concept for it to occur (Walker & Avant, 2011). Throughout the literature, an external or internal stimulant was found to occur repeatedly prior to the development of a certain behavior. In this case, the antecedent, which is the external or internal stimulant, happens as a prerequisite to the trigger. Another antecedent to trigger is the presence of ability and motivation. An individual must have the means and desire to perform a behavior.

Consequences are the result of a trigger taking place. The consequences of a trigger can be positive or negative. For example, an individual with asthma comes into contact with an environmental irritant. The irritant would exacerbate the individual’s asthma symptoms and thus would be negative. On the other hand, when an individual experiences a trigger that results in definitive action toward healthy lifestyle habits, the trigger is considered to be positive. In this case, the consequence is similar to a fight or flight response, or action toward a health outcome. However, the outcomes of trigger alone are inadequate to explain human behavior without some understanding of the individual’s motivation to actually take action.

EMPIRICAL REFERENTS

According to Walker and Avant (2011), empirical referents are a way to measure the concept under discussion and apply it to the real world. Throughout the literature, methods used to measure the presence of triggers mainly consisted of questionnaires specific to the condition under study (Andress-Rothrock et al., 2010; Chatterjee, Sherwin, & Jain, 2013; Ritz, Steptoe, Bobb, Harris, & Edwards, 2006; Visram et al., 2009). Ritz et al. (2006) developed the Asthma Trigger Inventory (ATI) to measure categories of patients’ self-reported asthma triggers. This questionnaire helped in initial screening of patients to determine the most relevant triggers for each individual, which allowed for evaluation of the impact of such triggers as well as the opportunity to address them in a specific way.

Triggers are unique to each individual. Thus, a questionnaire to determine the main categories of triggers that lead an individual to take action and change her or his behavior would be useful to effect behavior change.

MODEL CASE

According to Walker and Avant (2011), a model case is one in which all the defining attributes of the concept are exhibited. The following is a model case for trigger:

Janae is a registered nurse who is overweight and has recently been diagnosed with hypertension. One day, she is caring for a patient in the emergency room who has had a heart attack. Janae sees the pain and suffering in the patient’s face

*The firearm
industry uses
the word trigger
to describe the
lever on a gun
that the user
pulls to initiate an
action; this overt,
conscious, and
irrevocable action
provides a clear
mental image.*

and begins to think about her recent diagnosis of hypertension. As Janae reflects on this situation, she is determined not to let this happen to her. Once outside the room, she stomps her foot and declares, "That is not going to happen to me!"

After Janae finishes her shift, she drives to the gym to exercise. Instead of drinking alcohol after her shift, something she usually does, Janae chooses to drink water and eat a healthy meal.

This scenario exemplifies a model case because it includes all of the defining attributes of trigger. Caring for a patient experiencing complications of high blood pressure served as a trigger for Janae to take *unique*, definitive *action*. Upon reflection on the situation, Janae decided to change her behavior (*internal spark*).

Triggers can lead to both healthy and unhealthy behaviors. A study conducted by Kempes, Tiggemann, and Hollitt (2014) investigated the impact of exposure to televised food advertising and found that advertisements may activate thoughts about eating, triggering one's motivation to eat. Conversely, a vibration or alarm from wearable technology could be a reminder to stand, exercise, or go to sleep, thus triggering healthy behaviors that may decrease the risk of obesity.

CONCLUSION

The adoption of healthy behaviors is vital to reduce the burden of chronic illness. An important piece of this conversation must include identifying specific actions or cues that serve as motivational triggers. Despite interest in the obesity epidemic and numerous interventions to decrease its occurrence, overweight and obesity are continuing to increase (Dalle Grave et al., 2013). The increasing trend is detrimental to individuals, families, and the health care system alike. Determining the internal or external stimuli that trigger individuals to make a change and to take action is imperative to decrease the obesity epidemic.

Triggers can change over time (Peterson et al., 2012). According to Fogg (2009), a behavior will not occur without an appropriate trigger. With the increase in obesity, a method for sustained behavior change is necessary. The harmonious blending of motivational currents with positive behavior triggers will be a huge step in the control and treatment of behavior-based health challenges. Nurses should lead the way in this health care revolution based on positive individual changes to ensure that a happy, healthy lifestyle becomes the norm rather than the exception.

REFERENCES

- Andress-Rothrock, D., King, W., & Rothrock, J. (2010). An analysis of migraine triggers in a clinic-based population. *Headache*, 50(8), 1366–1370.
- Chambers, J. A., & Swanson, V. (2012). Stories of weight management: Factors associated with successful and unsuccessful weight maintenance. *British Journal of Health Psychology*, 17(2), 223–243.
- Chatterjee, K., Sherwin, H., & Jain, J. (2013). Triggers for changes in cycling: The role of life events and modifications to the external environment. *Journal of Transport Geography*, 30, 183–193. <http://dx.doi.org/10.1016/j.jtrangeo.2013.02.007>
- Dalle Grave, R., Centis, E., Marzocchi, R., El Ghoch, M., & Marchesini, G. (2013). Major factors for facilitating change in behavioral strategies to reduce obesity. *Psychology Research and Behavior Management*, 6, 101–110. <http://dx.doi.org/10.2147/PRBM.S40460>
- De Wet, C., & Bowie, P. (2011). Screening electronic patient records to detect preventable harm: A trigger tool for primary care. *Quality in Primary Care*, 19(2), 115–125.

The outcomes
of triggers alone
are inadequate
to explain
human behavior
without some
understanding of
the individual's
motivation to
actually take
action.

- Eaton, D. K., Kann, L., Okoro, C. A., & Collins, J. (2007). Selected health status indicators and behaviors of young adults, United States—2003. *American Journal of Health Education*, 38(2), 66–75.
- Fogg, B. J. (2009). *A behavioral model for persuasive design*. Retrieved from http://www.bjfogg.com/fbm_files/page4_1.pdf
- Fogg, B. J. (2011). *BJ Fogg's behavior model*. Retrieved from <http://behaviormodel.org.html>
- Jago, R., Nicklas, T., Yang, S. J., Baranowski, T., Zakeri, I., & Berenson, G. S. (2004). Physical activity and health enhancing dietary behaviors in young adults: Bogalusa Heart Study. *Preventive Medicine*, 41, 194–202. <http://dx.doi.org/10.1016/j.ypmed.2004.09.045>
- Kemps, E., Tiggemann, M., & Hollitt, S. (2014). Exposure to television food advertising primes food-related cognitions and triggers motivation to eat. *Psychology & Health*, 29(10), 1192–1205. <http://dx.doi.org/10.1080/08870446.2014.918267>
- Kungu, K., & Machtmes, K. (2009). Lifelong learning: Looking at triggers for adult learning. *International Journal of Learning*, 16(7), 501–511.
- Lim, D., Melucci, J., Rizer, M. K., Prier, B. E., & Weber, R. J. (2016). Detection of adverse drug events using an electronic trigger tool. *American Journal of Health-System Pharmacy*, 73, S112–S120. <http://dx.doi.org/10.2146/ajhp150481>
- Mohindra, N. A., Nicklas, T. A., O'neil, C. E., Yang, S.-J. T., & Berenson, G. S. (2009). Eating patterns and overweight status in young adults: The Bogalusa Heart Study. *International Journal of Food Sciences & Nutrition*, 60, 14–25.
- Nelson, M. C., Kocos, R., Lytle, L. A., & Perry, C. L. (2009). Understanding the perceived determinants of weight-related behaviors in late adolescence: A qualitative analysis among college youth. *Journal of Nutrition Education and Behavior*, 41(4), 287–292. <http://dx.doi.org/10.1016/j.jneb.2008.05.005>
- Nelson, M. C., Lust, K., Story, M., & Ehlinger, E. (2008). Credit card debt, stress and key health risk behaviors among college students. *American Journal of Health Promotion*, 22(6), 400–407.
- O'Toole, J., Solan, L., Burkhardt, M., & Klein, M. (2013). Watch and learn: An innovative video trigger curriculum to increase resident screening for social determinants of health. *Clinical Pediatrics*, 52(4), 344–350.
- Peterson, M. G., Gaeta, T. J., Birkhahn, R. H., Fernández, J. L., & Mancuso, C. A. (2012). History of symptom triggers in patients presenting to the emergency department for asthma. *The Journal of Asthma*, 49(6), 629–636.
- Quick, V., Wall, M., Larson, N., Haines, J., & Neumark-Sztainer, D. (2013). Personal, behavioral and socio-environmental predictors of overweight incidence in young adults: 10-Yr longitudinal findings. *International Journal of Behavioral Nutrition and Physical Activity*, 10(1), 37–49. <http://dx.doi.org/10.1186/1479-5868-10-37>
- Ritz, T., Steptoe, A., Bobb, C., Harris, A., & Edwards, M. (2006). The Asthma Trigger Inventory: Validation of a questionnaire for perceived triggers of asthma. *Psychosomatic Medicine*, 68(6), 956–965.
- Trigger. (n.d.). In *Merriam-Webster's online dictionary* (11th ed.). Retrieved from <https://www.merriam-webster.com/dictionary/trigger>
- Visram, S., Crosland, A., & Cording, H. (2009). Triggers for weight gain and loss among participants in a primary care-based intervention. *British Journal of Community Nursing*, 14(11), 495–501.
- Walker, L. O., & Avant, K. C. (2011). *Strategies for theory construction in nursing* (5th ed.). Upper Saddle River, NJ: Prentice Hall.
- Wong, B., Dyal, S., Etchells, E., Knowles, S., Gerard, L., Diamantouros, A., . . . Shojania, K. (2015). Application of a trigger tool in near real time to inform quality improvement activities: A prospective study in a general medicine ward. *BMJ Quality & Safety*, 24(4), 272–281.

Correspondence regarding this article should be directed to Christine S. Gipson, PhD, RN, CNE, at cgipson@uttyler.edu